

Multifactor mechanism of SCENAR-analgesia

A.V. Tarakanov, E.G. Los

Reflexology Journal, Issue No: 3 (7), 2005 Moscow, Russia

Department of "Faculty of Qualification and Improvement in A&E Departments" and "Professional Training of Specialists" at State Medical University of Rostov

Original in Russian:

А. В. Тараканов, Е. Г. Лось: Многофакторный механизм СКЭНАР-Анальгезии; Научно-практический Журнал "РЕФЛЕКСОЛОГИЯ" № 3 (7), 2005

Abstract:

The complex effect of SCENAR-therapy is demonstrated in the treatment for various conditions associated with acute pain in emergency situations.

The result of monotherapy of pain:

- SCENAR had an analgesic effect for patients with traumas (n=20, 60%);
- Good analgesia was also achieved with SCENAR in patients with a background of stenocardia and normalisation of functions of the heart (n=26, 66%) and
- in hypertensive crises (n=153, 88-91% for cephalgia, and cardialgia respectively).

Introduction

The issue of protecting patients against pain has an enormous humanistic importance. Pain is a signal of danger, a symptom of many diseases, is the main reason for patients to suffer, and a basic pathophysiological factor for developing shock.

When a patient faces aversive factors in the environment, adaptive mechanisms are started with the aim for the organism to survive in the new conditions. This is mainly done by the activation of the endogenous anti-nociception system, one of the general biologic mechanisms. It is known that the mechanisms of endogenous anti-nociception are mediated by opiate and non-opiate systems of the brain. The mechanisms of anti-nociception have been subject of numerous comprehensive works, including acupuncture, electropuncture, and transcutaneous neurostimulation. [2,3,8,11]

It is known that pharmaceutical methods of analgesia are imitations of the functions of the endogenous system to protect against pain. They are important when performing mass analgesia.

Starting the endogenous anti-nociception system is, however, optimal when methods of reflex therapy are used, because this method is more physiologic. The complexity of the methods, their invasiveness, difficulty to be used in extreme circumstances, lack of reliable and inexpensive equipment, and a lack of trained specialists are the factors that obstruct a wide introduction of reflexological methods for pain relief into main-stream medicine.

Methods of transcutaneous bio-regulated, low-frequency impulse-electrotherapy have been used recently for non-pharmaceutical treatment [5,6,7], particularly the Self-Controlled Energo Neuro Adaptive Regulator (SCENAR). This method is based on the effects of a bipolar impulse of low frequency without a constant (DC) component. Characteristics which distinguish SCENAR among other electric equipment for transcutaneous stimulation are:

- the optimal form of the impulse and the force of its effect;
- the body almost does not have to adapt to the SCENAR impact due to its biotechnological feed-back;
- non-damaging mode of impact;
- high level abruption of the first part of the affecting signal with neutralization of the accommodation effect;
- many researches have been performed to examine this method of treatment;
- during the treatment, an expert evaluation of the dermal condition can be used to optimize the area of impact;
- optimal construction of the device enables it to work in various conditions [6, 7].

Applying SCENAR-therapy is an issue of huge interest due to its availability, rapid positive effect, simplicity of treatment and steady result.

Aim of the Research

Examining possibilities to perform SCENAR-therapy at various pathological processes accompanied by acute pain in emergency situations.

Materials and Methods

Mobilizing the adaptive systems of the organism with SCENAR in situations of acute damage to their functions and to achieve an analgesic effect were the reason to use the SCENAR in emergency situations. More than 40 doctors of accident & emergency departments in 13 towns in the districts of Rostov, Stavropol, Nizhni Novgorod and Adigeia participated in this research. More than 500 patients were treated. The coordination of the clinical tests according to a specially developed plan was performed by the “Faculty of Qualification Improvement for Emergency Departments” of the State Medical University in Rostov in conjunction with the department of clinical research of “RITM OKB” Development Bureau in Taganrog [the manufacturer of SCENAR devices]

Patients with acute pain due to three different pathological processes were randomly selected from the database:

- 1) traumatic injuries of limbs, soft tissue injuries (n=20);
- 2) steady stenocardia (n=26);
- 3) hyper-intensive crises (n=153).

SCENAR-therapy was performed according to instructions [12]:

- Local Zones including “anti-pain” points;
- If needed general zone treatments could be added:
 - “three paths on the backbone and six points on the face” and
 - “collar area, forehead, adrenals” (Collar Zone).
- If there was no possibility to affect the injured organ, the device was applied to the “symmetric areas”.

The skin was treated using a CHENS-SCENAR device without digital methods in mode F1 (constant or manual mode). The therapeutic procedure was performed only once by an emergency department doctor, as the duration of the procedure was 5-30 minutes long, depending on the effect. When the analgesia was ineffective, the doctor could inject analgesic medicine. To evaluate the effect of the analgesia in conditions of emergency, a visual analog scale (VAS) was used, where 0cm = no pain, 10cm = maximum pain.

The use of the protocol **“three paths of the spine and six points of the face”** was performed on the grounds of the following:

- Processing path 2 and 3, Shu-points of the back are included in the area of the stimulation with SCENAR (points of consent) [4, 10, 11], which are located on the Bladder Meridian (BL).
- Path 1 covers the unpaired rear middle meridian, the Governing Vessel (GV).
 - When areas containing points GV 1- GV 4 are processed a therapeutic effect is made on this meridian aimed at nervous system diseases, intestinal diseases, diseases of the urino-genital system.
 - When areas containing points GV 5- GV 8 are processed a therapeutic effect is made on this meridian aimed at nervous system diseases, diseases of gastric and intestinal system.

- The area of points from GV 9 to GV 14 – regulate nervous system disorders, pulmonary diseases, fever.

Processing six points on the face enables to relay information through all the three rami of the trigeminal nerve. On the other hand, biologically active points of the meridians are located in these areas. At least point BL 2 (Bladder Meridian), points St 2 and St 3 (Stomach Meridian), Li 20 (Large intestine Meridian), and point St 4 (Stomach Meridian) are included in the area of impact by this SCENAR protocol.

The method **“collar, forehead, adrenals (Collar Zone)”** is also related to the areas of general treatment. This protocol is useful as it covers large reflexogenic areas and biologically active points - as the experience of SCENAR-therapy shows [5, 12]. When the indicated protocol is used and when the zones are processed “according to instructions”, the device may stick to the skin. In this case, the electrode is not lifted off the skin but is held until further movement becomes possible. If the device does not stick, then other kinds of so-called asymmetries can appear (hyperaemia, change of the characteristics of the sound, different sensibility etc.). These dermal areas are worked on additionally until initial characteristics are changed.

Results and Discussion

1 Pain at traumas. Analgesia with the use of SCENAR was performed at small and medium traumas: lower limbs fracture (n=5), injuries of soft tissues of the limbs and thorax (without damaging respiratory functions) (n=9), I and II degree burns of face and abdominal wall (n=2), Cuts and bites (n=4). There were nine men and 11 women with age from 11 to 82 (average age – 40.7 ± 3.9). The patients were treated using general means of treatment: splinting, immobilization, non-adhesive bandages in cases of burn etc. The evaluation of the pain was performed using VAS before the analgesia with SCENAR, immediately after the procedure, and in the 10th and 20th minute. In the same intervals, other complaints and symptoms were assessed, as well as the parameters of haemodynamic. The data are shown on fig.1.




Fig.1. **A** - Dynamics of the analgesic effect of SCENAR during therapy of traumatic pains according to VAS. **B** – Dynamics of systolic and diastolic blood pressure and pulse frequency of patients with traumatic pains. Reliability referring to data before SCENAR: * - $P < 0.05$; ***- $P < 0.001$;

A distinct reliable analgesic effect developed immediately after the procedure, almost without further increase by the 20th minute. Observations showed that the analgesic effect increases by the 50th – 60th minute. Because the emergency staff cannot follow the efficiency of the procedure for a long period of time, in several cases the doctor made the decision to inject additionally non-opiate analgesic medicine (analgin, ketorolac or ketonal). Only 8 out of 20 patients were injected analgesic medicines. Natural skepticism and caution should be noticed when doctors tested the new method.

We regard as an important factor the steadiness of the values of blood pressure. Reliable decrease of the average frequency of cardiac contractions from 85 to 61 beats per minute indirectly shows sympathetic tonus. A reliable change of the respiratory frequency was not noticed and it was within normal limits. Decrease of other symptoms and complaints was

noticed along with the analgesic effect, such as dizziness, nausea, excitedness and sensation of inhibition.

Independent analgesic effect developing on the background of steady parameters of blood pressure and decreasing of symptoms accompanying the traumas is noticed during the process of analgesia using SCENAR for patients with small and medium traumas.

2 Pain at steady stenocardia tension of I-III functional class.

26 patients with ischemic cardiac disease, crises of pain at steady tension stenocardia of different functional classes were treated with SCENAR, as the patients had previous idiosyncrasy to nitrates or lack of efficiency of the nitrates. There were 14 men (average age – 63,1,7±2,5 years old), 12 women (average age - 68.4±3.7 years old). 7 patients had arterial hypertonia as accompanying disease, and 3 of them had osteochondrosis of the backbone. This category of patients was additionally treated with SCENAR as the dermal area of the pericardial channel was processed in the lower part of right forearm and the painful area of skin on the thorax.





Fig.2. **A** – Dynamics of the complete analgesic effect and number of residual pains (%) during analgesia of patients with attacks of stenocardia. **B** - Dynamics of analgesia of patients with residual pains during SCENAR-therapy, according to VAS. Reliability referring to SCENAR data: * - P < 0.05; ** - P < 0.02; *** - P < 0.001.



As can be seen in fig.2 (A) immediately after SCENAR-treatment the pain completely disappeared for 46 % of the patients and for 65% of the patients – 20 minutes later. 35% of the patients felt residual pain in the 20th minute after the treatment, but the intensity of the pain was much less, which can be seen in fig 2 (B). These patients, among whom there were mainly patients with accompanying diseases, were offered additional therapy with included symptomatic medicines and non-opiate analgesics. Changes of parameters of blood pressure and pulse statistically did not differ for the category of patients with additional treatment and patients with SCENAR-therapy.

This way, analgesia using SCENAR for algic crises at steady tension stenocardia causes sufficient independent analgesic effect, which does not depend on parameters of general pressure. The extent of its correlation with coronary varicose and the endogenous mechanisms which SCENAR starts in order to stop the pain, can be defined in hospital conditions and additional tests.

3. Cardialgia and cephalgia at hypertensive crises.

During SCENAR-therapy for patients with neurovegetative forms of hypertensive crises normalization of the blood pressure was an aim, not only the analgesic effect. A total of 153 patients were helped at their homes on emergency call-outs. To analyze the analgesic effect we divided the patients into 2 groups: with cephalgia (n=137, which forms 90% of the patients; among them 71 patients – 52% had cardialgia); separately data of patients with cardialgia were analyzed (n=78, which is 51% of the patients). Patients of this category were treated with SCENAR: in the collar zone, area of pericardial meridian, in the lower part of the right forearm, and dermal projections of maximal pain.





Fig.3. A – Dynamics of the complete analgesic effect and number of residual pains (%) during analgesia of patients with cardialgia at hypertensive crisis. **B** - Dynamics of analgesia of patients with residual pains during SCENAR-therapy, according to VAS. Reliability referring to SCENAR data: * - $P < 0.05$; ** - $P < 0.02$; *** - $P < 0.001$.



In fig. 3 can be seen that immediately after the procedure cardiac pains disappeared for 61% of the patients, and for 91% by the 30th minute. Nevertheless, residual painful sensations were registered by the 30th minute only for 9% of the patients. Their extent was much lower, as at stenocardia, which could be seen during evaluation using visual analog scale (fig. 3 B). This category of patients at times needed an additional injection of non-opiate analgesic.

In fig.4, the effect of the SCENAR device can be seen for cephalgia and hypertensive crises at pre-hospital stage of treatment. Fig.4 A shows reliable and gradual elimination of pain. After the procedure, the headache disappeared for 43%, and for 88% of patients - by the 30th minute.

Residual pains by the 30th minute were registered only for 12% of the patients. Their extent, according to VAS, was much lower than before the procedure (Fig.4 B). Unidirectional elimination of pain in cardiac area and headaches was noticed during elimination of hypertensive crisis. We analyzed the changes of parameters of blood pressure during SCENAR-therapy.

Some patients took hypotensive medicines before the emergency staff arrived, which dramatically changes the efficiency of SCENAR. Naturally there was certain skepticism: was it SCENAR that affected or was it previously taken medicines?





Fig.4. A – Dynamics of the complete analgesic effect and number of residual pains (%) during analgesia of patients with cephalgia at hypertensive crisis. **B** - Dynamics of analgesia of patients with residual pains during SCENAR-therapy, according to VAS. Reliability referring to SCENAR data: * - P < 0.05; ** - P < 0.02; *** - P < 0.001.



This is why we divided all the patients into 2 groups: one who took medicines and one who did not take medicines before they called the emergency and were treated with SCENAR. Among the patients who had taken hypotensive medicines before the emergency arrived (calcium antagonists, β -adrenoblockers, APF inhibitors etc) 30-90 minutes before that without any effect there were 80 people (58 women, 22 men; average age – 61.1 \pm 4.9 years old). The group of patients who did not take medicines before calling the emergency contained 73 people (52 women, 21 men; average age – 63.1 \pm 3.5 years old). The results are shown in fig.5.

As can be seen in the left part of the figure, the parameters of blood pressure for both groups did not differ before and after SCENAR-treatment, on the background of general decrease of blood pressure. 30 minutes before the end of the procedure the systolic blood pressure decreased 11-16%, diastolic – 11-9%. This speed of decrease of blood pressure in the first 30 minutes is effective as cardialgia and cephalgia are eliminated it is safe for all ages. A typical effect of the SCENAR-therapy is the primary elimination of the complaint of the symptom and

then normalization of objective parameters. Vegetative corrective effect of SCENAR-therapy should be noticed as hyperhydrosis, nausea, vomiting and excitedness is eliminated. Further observation of some patients and the experience of elimination crises in hospital conditions show that average blood pressure decreases 15-25% within 1 hour, which does not lead to ischemia of targeted organs. The reliable decrease of frequency of cardiac contractions with 9% could be noticed when eliminating crises without previous use of medicines (fig.5b)

A





Fig.5. A – Dynamics of parameters of blood pressure during SCENAR-therapy of hypertensive crises in the group of patients who had taken and who had not taken hypotensive medicines before the emergency staff arrived.

B - Dynamics of frequency of cardiac contractions in the same groups. Reliability referring to the data before SCENAR: * - $P < 0.05$

B






Fig.6. Dynamics of correlation coefficient depending on the decrease of systolic blood pressure and decrease of painful sensibility at cephalgia and cardialgia, according to VAS.

If during the process of analgesia, the analgesic effect develops independently for patients with traumas, then during attack of pain at steady stenocardia and particularly hypertensive crises, the analgesia probably depends mostly on recovering of functions and decrease of ischemia: the decrease of system blood circulation and normalization of regional blood circulation including the coronary one.

We analysed the dynamics of correlation coefficient depending on the decrease of systolic blood pressure and the extent of headache and cardiac ache, according to visual analog scale. The data is represented in fig.6. As can be seen in this figure the correlation coefficient increases simultaneously with the decrease of the systolic blood pressure and decrease of values of extent of cardialgia and cephalgia, according to VAS. If in the beginning of the crisis the extent of the pain did not depend on the values of blood pressure, then SCENAR-therapy led to an increase of the correlation to 0.41 and 0.48. Similar dynamics of close relation and its lack in the beginning of the crisis probably shows simultaneously developing effects of SCENAR: analgesic effect and the effect, which normalizes the functional system of blood pressure regulation. This way, the elimination of neurovegetative form of hypertensive crises using SCENAR regardless of preliminary therapy leads to gradual decrease of parameters of blood pressure and frequency of cardiac contractions. A distinct analgesic effect of cephalgia and cardialgia is noticed, which probably has a mixed nature.

During the process of elimination of acute pain with different origins, three elements should be present in the algorithm: possible elimination of reason causing pain; affecting the sensation of pain - perception; possible prophylactics of pathophysiological damages, which pain causes or will cause.

As the experience of work with SCENAR shows, during the treatment of acute and chronic painful syndromes, the data presented in this article show the multicomponentness of the suggested therapy. Non-medication-method for acute pain relief starts the endogenous antinociception and other mechanisms of recovering damaged functional systems. The analysis of the data obtained in traumatology and cardiology shows that the share of perceptual components of SCENAR analgesia is different depending on nosology, the level and the volume of the damage. It is not reasonable to rely on only one single method of analgesia and it is also dangerous for the patient, especially in case of emergency.

Undoubtedly, the morphologic substrate of the analgesic effect of SCENAR-therapy are the antinociceptional systems of brain, starting from the different areas of skin. SCENAR works with the biologically active points, with their accumulation and the surrounding tissues – this is a zonal device. Due to its technical solution, it “finds” these points independently. As shown for acupuncture analgesia [8], in opioidergic, serotonergic, catecholaminergic, cholinergic, gamkergic and possibly other mechanisms take part in its realization.

Some authors regard the transcutaneous electro-stimulation as the most accessible and the simplest “hyperstimulation analgesia” to be realized. If we proceed from the theory of Melzack and Wall (1965) the activity in the thin non-myelinated C-fibers located in the substantia gelatinosa of the dorsal horns of the spinal cord “opens the gates” to the further pass of nervous impulses bearing the information of pain. The role of the thick fibers consists in the ability to “close the gates”. One of the hypotheses of the analgesic effect of transcutaneous neurostimulation consists of the fact that the procedure leads to activation of thick myelinated fibers with subsequent “locking the painful entrance”. The analgesic effect during the so-called dynamic neurostimulation is proved to be naloxone-dependent [9].

Using almost the same methods of SCENAR-therapy and taking under consideration the principles of their application [5,6,12], we obtained unidirectional results: analgesic effect and final positive sanogenic result connected with normalization of the changed functional system. Taking under consideration the limited volume of such a publication we can mention that similar unidirectional results were obtained during the process of therapy of acute myocardial infarction, burns, bronchial asthma of adults and infants etc.

The simplicity of use of SCENAR allows us to recommend the use of this kind of analgesia in any conditions and if needed, by people without medical education.

Conclusions

1. The SCENAR-therapy (self-controlled, energy neuro-adaptive regulator) is an effective, safe and multifactor non-pharmaceutical method of analgesia during treatment of different conditions accompanied by acute pain in cases of emergency.
2. During the process of analgesia using SCENAR the patients with small and medium trauma (n=20), an independent analgesic effect is noticed (60%) developing on the background of steady parameters of blood pressure and decrease the symptoms of the accompanying traumas.
3. The analgesia of painful attacks from ischemic disease and steady tension stenocardia (n=26) using SCENAR, causes a sufficient independent analgesic effect (66%), which does not depend on the parameters of general pressure.
4. The elimination of neuro-vegetative form of hypertensive crises (n=153) using SCENAR, regardless of preliminary medical therapy leads to gradual reliable decrease of the frequency of cardiac contractions and decrease the parameters of blood pressure by the 20th minute after the procedure; by the 30th minute a distinct analgesia of cephalgia and cardialgia is noticed (88-91% respectively).

Bibliography:

1. Abakyan G.N. Reflexotherapy of diseases of the nervous system. M., 1999. - 297p.
2. Barashkov G.N. Reflexotherapy of pain. - M.: НЦМ "Универсимед", publishing house "ТМ - ОКО ", 1995. - 263p.
3. Vasilenko A.M. Acupuncture and reflexology: the evolution of methodology and metatheory. - Taganrog. 1998. - 110s.
4. Vogralik VG, Vogalik M.V. Acupuncture. The foundations of the traditional eastern Reflexodiagnosics and acupuncture adaptive-energizing therapy: Tsi-Gong. - M.: GOU VUNMTS of the Russian Federation, 2001. - 336s.
5. Gorfinkel Yu.V. - Theoretical and practical basis for improving Effectiveness of SCENAR-therapy. In the collection of articles "SCENAR-therapy and SCENAR-Espertiza ", issue 2, Taganrog, 1999. - P. 16-18
6. Grinberg Ya.Z. To the question on the basis of the effectiveness of SCENAR-therapy / SCENAR - therapy and SCENAR examination. - Taganrog 1997. - Issue 3. - p.16-22.
7. Grinberg Ya.Z. The concept of electrotherapy / SCENAR-therapy and SCENAR-expertise. Taganrog, 1999. - issue 5 - p.16-13
8. Ignatov Yu.D., Kachan AT, Vasilyev Yu.N. Acupuncture Analgesia: Experimentally-clinical aspects. L. Meditsina, 1990. - 256s.
9. Kukushkin ML, Meizerov EE, Grafova V.N. Development features analgesic effect in transdermal dynamic electroneurostimulation // Bulletin of Experimental Biology and Medicine. - 2003. - №3. - P.265-267
10. Luvsan Gaava. Essays on the method of eastern reflexotherapy. Novosibirsk: Science, 1991. - 432p.
11. Machareth EL, Korkushko A.O. Basics of electro and acupuncture - Kiev: Health, 1993. - 391s.
12. Revenko AN, Tarakanov A.V. Etc. Instructions for the use of percutaneous Electroneurostimulator with bio-inverse communication of individually-dosed Influence on the reflex zones. CHENS-SCENAR. - Taganrog, 2004. 70s

Translated from Russian to English Language by: Mrs. Veselina Petkovap August 2008, Mladost 329, Sofia 1712, Bulgariap <http://syntagma.dir.bg>

Translation revised by Dr. med. Jorg Prinz, Energy Health Clinic, Whakatane New Zealand 2017